



## AMERICAN SOCIETY OF ANESTHESIOLOGISTS CLASSIFICATION

- I. Healthy Patient
- II. Mild systemic disease - no functional limitations
- III. Severe systemic disease - definite functional limitation
- IV. Severe systemic disease that is a constant threat to life
- V. Moribund patient - not expected to survive 24 hours with or without operation
- E. Emergency

### ALDRETE SCORE:


A score of 10 is required for discharge from post procedure area. For discharge with a score less than 10, a Doctor's specific order is required.

## PAIN SCALES


### FLACC Scale

Categories	Scoring		
	0	1	2
Face	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested	Frequent to constant quivering chin clench jaw
Legs	Normal position or relaxed	Uneasy, restless, tense	Kicking, or legs drawn up
Activity	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid or jerking
Cry	No cry (awake or asleep)	Moans or whimpers; occasional complaint	Crying steadily, screams or sobs, frequent complaints
Consolability	Content, relaxed	Reassured by occasional touching, hugging or being talked to, distractable	Difficult to console or comfort
Each of the five categories (F) Face; (L) Legs; (A) Activity, (C) Cry; (C) Consolability is scored from 0-2, which results in a total score between zero and ten.			


### Visual Analog Scale




### Alternative Assessment Tool Non-communicative




## MALLAMPATI CLASSIFICATION




CLASS I



CLASS II



CLASS III



CLASS IV